


Role of pulsed electromagnetic field on enzyme activity, germination, plant growth and yield of durum wheat

Katsenios. Nikolaos, Bilalis Dimitrios, Efthimiadou Aspasia, Aivalakis Georgios, Nikolopoulou Aimilia-Eleni, Karkanis Anestis, Travlos Ilias



www.elsevier.com/locate/bab

PII: S1878-8181(16)30049-4
DOI: <http://dx.doi.org/10.1016/j.bcab.2016.03.010>
Reference: BCAB378

To appear in: *Biocatalysis and Agricultural Biotechnology*

Received date: 2 February 2016
Revised date: 27 March 2016
Accepted date: 29 March 2016

Cite this article as: Katsenios. Nikolaos, Bilalis Dimitrios, Efthimiadou Aspasia Aivalakis Georgios, Nikolopoulou Aimilia-Eleni, Karkanis Anestis and Travlos Ilias, Role of pulsed electromagnetic field on enzyme activity, germination, plant growth and yield of durum wheat, *Biocatalysis and Agricultural Biotechnology* <http://dx.doi.org/10.1016/j.bcab.2016.03.010>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

**Role of pulsed electromagnetic field on enzyme activity, germination, plant growth and
yield of durum wheat**

Katsenios Nikolaos¹, Bilalis Dimitrios¹, Efthimiadou Aspasia², Aivalakis Georgios³,
Nikolopoulou Aimilia-Eleni³, Karkanis Anestis⁴, Travlos Ilias¹

¹Laboratory of Crop Production, Agricultural University of Athens, Iera Odos 75, 11855
Athens, Greece

²Open University of Cyprus, P.O. Box 24801, 1304 Nicosia, Cyprus

³Laboratory of Plant Physiology, Agricultural University of Athens, Iera Odos 75, 11855
Athens, Greece

⁴Department of Agriculture Crop Production and Rural Environment, University of Thessaly,
Fytokou Street, N. Ionia, 38466 Magnisia, Greece

Abstract

Researchers have focused their efforts on the use of magnetic field as a pre-sowing method, as it is an inexpensive, environmentally friendly technique. This study provides a holistic approach of an agricultural cultivation that can lead to the comprehension of the exact mechanism of magnetic field effect on plant and lead to the appropriate application of magnetic fields. Pulsed electromagnetic field was used for 0, 15, 30 and 45 minutes as a pre-sowing treatment in durum wheat seeds in a field experiment for three years. The experiment followed a completely randomized design, with four treatments (Control, MF-15, MF-30 and MF-45), two cultivars and three replications. The aim of this study was to determine the effect of magnetic field exposure on durum wheat seeds, covering a complete range of agronomic characteristics. The results obtained in this three-year experiment showed a positive impact of pulsed electromagnetic field, in durum wheat cultivation. Magnetic field

has been found to enhance germination, tillering, dry weight, leaf area, chlorophyll content, photosynthetic rate, transpiration rate, stomatal conductance and yield of two durum wheat cultivars. Duration of 30 and 45 minutes of pre-sowing magnetic field treatment gave the best results. Regarding yield measurement, all magnetic field treatments (MF-15, MF-30 and MF-45) gave statistically significantly higher values compared to control. In addition, α -amylase activity measurements showed that magnetic field affects enzymes and this could possibly explain the improved germination.

Keywords: magnetic field, α -amylase, germination, durum wheat, seed priming, pre-sowing

INTRODUCTION

Investigation of magnetic field's influence on plant growth has started to develop rapidly during the last decades. Encouraging results indicated that magnetic field could enhance some plant functions. The use of magnetic field on seeds as a priming technique is getting more and more familiar among researchers (Bhardwaj et al., 2012).

The existence of earth's magnetic field is known to mankind since ancient times, through the discovery of the ability of the mineral magnetite to be oriented in the direction North - South when freely suspended above the earth (Stavroulakis, 2003). But despite the fact that it has always been one of the key features of the environment, until recently the study of the effect of the magnetic field in agronomic science was absent. Each region is characterized by a number of abiotic factors (water, air, soil, temperature) including the magnetic field. However, for many years, the magnetic field of earth constituted as an invariable parameter of the environment that was not taken into account in plant studies (Katsenios, 2013).

The father of bio-electromagnetism is Hippocrates, who first tried to cure breast cancer by exposure of the sun's electromagnetic radiation (therapy by the sun's rays). About 2,000 years later, during the 18th century, Luigi Galvani tried to treat tumors, aneurysms and hemorrhages by applying electricity to tissues. In 1840, Recamier and Pravaz provided a method of destroying the cancer cells in the uterus through the use of electricity, which soon became common practice (Stavroulakis, 2003).

Different types of plant materials (seeds, seedlings, young plants and cuttings) have been used. These plant materials have been treated with different types of magnetic field. (Flórez et al., 2007; Hajnorouzi et al., 2011). More particularly, in order to study the effects on plants, various types of magnetic fields have been used (static, electromagnetic, pulsed electromagnetic), in different intensities and exposure duration (Poinapen et al., 2013, Aguilar et al., 2009, Bilalis et al., 2012a, Radhakrishnan and Kumari, 2012). Regarding exposure time, it varies from 15 seconds (Muszinski et al., 2009) to 24 hours (Martinez et al., 2009). Many experiments have been done with cereal grains (Torres et al., 2008; Vashisth & Nagarajan, 2008; Vashisth & Nagarajan, 2010), legumes (Podlesny et al., 2004; Podlesny et al., 2005; De Souza et al., 2006, Odhiambo et al., 2009;), and perennials (Çelik et al., 2008; Dardeniz et al., 2006; Dhawi & Al-Khayri, 2009).

Investigations in various plant species showed a positive effect of the magnetic field on seed germination (Bhardwaj et al., 2012), plant development in the early stages (De souza et al., 2014) and ultimately yield (Vashisth et al., 2013). Magnetic field has been found to improve the growth of plants, such as tomatoes (De Souza et al., 2006), the germination and the early stages of growth in plants such as sunflower (Vashisth & Nagarajan, 2010) and soybean (Radhakrishnan and Kumari, 2012). Moreover, researchers report the increase of the root system in young maize seedlings (Muraji et al., 1998). Animal experiments have shown that

short duration PEMF seem to facilitate and improve the quality of skin wound healing in rats (Athanasίου et al., 2007).

The enhancement of plant growth parameters is absolutely desirable in modern and organic agriculture, since it can be achieved through the use of an environmentally friendly method, which is also inexpensive. It is worth noting that many researchers have focused their efforts on the use of magnetic field as a pre-sowing technique, as it is an inexpensive, environmentally friendly technique, which can be applied with relative ease (Vashisth and Nagarajan, 2010).

Magnetic field treatment on seeds has been found to increase the activity of hydrolytic enzymes such as amylases in sunflower (Vashisth and Nagarajan, 2010). In cucumber, magnetic field treated seeds showed higher activity of β -amylase and finally increased the rate of germination compared to control (Bhardwaj et al., 2012). The increase of enzyme activity could be a primary positive effect of magnetic field treatment that subsequently leads to higher germination percentage, plant growth and yield.

The aim of this study was to determine the effect of magnetic field exposure on durum wheat seeds, covering a complete range of agronomic characteristics such as germination, tillering, dry weight, leaf area, chlorophyll content, photosynthetic rate, transpiration rate, stomatal conductance and yield. The activity of α -amylase was investigated to explain the increase of germination of durum wheat seeds. Our purpose was to have replicable results in a three-year period.

MATERIALS AND METHODS

Plant Material

A field experiment was established at the Agricultural University of Athens (Greece), for three consecutive cultivation years (2009/10, 2010/11 and 2011/12). Two durum wheat (Simeto by Thrakiki Sporoparagogiki and Grecale by NAGREF) varieties were used. Simeto has a high yield potential with thousand grain weight (TGW) at 36-39 g and hectoliter weight at 78-82 kg/hl. Grecale has a medium to high yield potential with thousand grain weight (TGW) at 40 g or more and hectoliter weight at 80 kg/hl.

Treatments

Four different exposure times of pulsed electromagnetic field were applied in both cultivars. The seeds were treated using a PAPIMI electromagnetic field generator for 15, 30 and 45 minutes before planting. Non treated seeds were used as control. The PAPIMI device is a pulsed electromagnetic field (PEMF) generator (PAPIMI model 600; Pulse Dynamics, Athens, Greece. Manufacturer characteristics: 35–80 J/pulse energy, 1×10^{-6} s wave duration, 35–80 $\times 10^6$ W wave power, amplitude of the order of 12.5 mT, rise time 0.1 ms, fall time 10 ms, repetitive frequency of 3 Hz.). The same device has been used in medical and agricultural studies (Athanasίου et al., 2007; Bilalis et al., 2012b; Katsenios, 2013; Milgram et al., 2004).

Experimental design

The experiment followed a completely randomized design, with four treatments (Control, MF-15, MF-30 and MF-45), two cultivars and three replications for three years. Seeds were treated for 15 minutes (MF-15), 30 minutes (MF-30) and 45 minutes (MF-45). Non treated seeds were used as control (0 minutes). Every replication was consisted of an area of 6 m². The quantity of seeds used was 16 g / m² for all cultivars.

Measurements and Observations

Germination (number of plants per 1 meter row) measurement took place 20 DAS, while **tillering** (plants per 1 meter row) took place 50 DAS. **Leaf area** (cm^2 per plant) and **stem dry weight** (g per plant) were destructive measurements and took place 150 DAS. Leaf area was measured by using an automatic leaf area meter (Delta-T Devices Ltd., Burrwell, Cambridge, UK). Stem dry weight was measured by a precision balance after the samples were oven dried at 70°C for three days in order to measure the weight in grams per plant. For the **chlorophyll** ($\mu\text{g}/\text{cm}^2$) measurement, a portable chlorophyll meter (SPAD) was used. The measurement was taken 130 DAS. A calibration curve has been created in order to convert the SPAD measurement to $\mu\text{g}/\text{cm}^2$ (Lichtenthler and Wellburn, 1983). Measurements of **photosynthetic rate** ($\mu\text{mol CO}_2\text{m}^{-2}\text{s}^{-1}$), **transpiration rate** ($\text{mmol H}_2\text{O m}^{-2}\text{s}^{-1}$) and **stomatal conductance** ($\text{mol m}^{-2}\text{s}^{-1}$) were undertaken between the hours of 10.30 and 14.30 on fully expanded leaves, with five measurements per treatment. Measurements were made using an LCi Leaf Chamber Analysis System (ADC, Bioscientific, Hoddesdon, UK). Physiology measurements were taken 115 DAS.

The activity of **α -amylase** has been recorded on Simeto variety, for a single year. A reference curve has been produced, using standard solutions and then we took three measurements, the third, the fourth and the fifth day after sowing. The activity of α -amylase was determined using the method described by Guglielminetti et al., 1995. Samples (0.2-0.5 g fresh weight) were extracted in 100 mM Hepes-KOH, pH 7.5, containing 1 mM EDTA, 5 mM MgCl_2 , 5 mM DTT, 10 mM NaHSO_4 . Extracts were centrifuged (13,000g, 15 min), the resulting pellets were washed with the extraction buffer and centrifuged again, and the resulting supernatants were combined and used for the enzymatic assays. Samples were assayed for the enzymatic activities at 25°C in 0.5-mL reaction mixtures using the following method. α -amylase (Doehlert et al., 1982): samples pretreated at 70°C in the presence of 3 mM CaCl_2 , to eliminate interferences from β -amylase were incubated with 2.5% (w/v) soluble starch;

activity of enzyme (1 unit) is defined as the amount of enzyme required to produce 1 pmol Glc min⁻¹.

Statistical analysis

The experimental data were analyzed using Statistica software (StatSoft, 1996), according to the completely randomized design. Analysis of variance (ANOVA) and comparisons of means were calculated using the least significant difference (LSD) test, at the 5% level of significance.

RESULTS

The use of pulsed electromagnetic field as a pre-sowing treatment was found to enhance durum wheat germination, tillering, dry weight, leaf area, chlorophyll content, photosynthetic rate, transpiration rate, stomatal conductance and yield. Moreover, it was found that magnetic field increased the activity of α -amylase, the third and the fourth day after sowing.

Germination and Tillering

Magnetic field improved germination (plants per row) of durum wheat seeds (Table 1, 2). An interaction of magnetic field and variety has been recorded, in the three-year field experiment. The highest number of plants per row has been measured at *Grecale-45* treatment (80.96), with statistically significant differences compared to all other treatments, except *Grecale-30* (80.55) and *Grecale-15* treatment (78.80). These treatments gave values with no statistically significant differences compared to *Simeto-45* (78.34). Moreover, *Simeto-30* (73.17) and *Grecale-0* (71.68) gave higher values with statistically significant differences compared to *Simeto-15* (69.10), which gave higher values with statistically significant differences compared to *Simeto-0* (60.05). Tillering measurement (plants per row)

showed statistically significant differences among treatments (Table 1, 2). An interaction of magnetic field and variety has been found, in the three-year field experiment. The highest number of tillering has been found at *Grecale-45* (129.16), followed by *Grecale-30* (123.86). Both treatments gave values with statistically significant differences compared to all other treatments, with statistically significant differences. *Grecale-15* (112.81) gave higher values with statistically significant differences compared to *Grecale-0* (104.62). Moreover, *Simeto-45* (96.62), *Simeto-30* (95.73) and *Simeto-15* (92.04) gave values that were statistically significant higher compared to the control (78.17), for significant level of 0.05. Regarding year, the *third one* (100.20) gave values that were statistically significant higher compared to *first* and *second* year.

Plant growth

Plant growth characteristics gave statistically significant differences (Table 3). An interaction of magnetic field and variety has been found, in the three-year field experiment (Table 1). The highest values in plant dry weight (g per plant) were measured at *Simeto-45* treatment (15.23) with statistically significant differences compared to all treatments except *Grecale-45* (15.11), *Grecale-30* (14.64), *Simeto-30* (14.05) and *Grecale-15* (14.02). Moreover, *Simeto-15* (13.81) gave values that were statistically significantly higher compared to *Simeto-0* (12.03) and *Grecale-0* (11.94), for significant level of 0.05. Regarding year, the *second* (14.82) and the *first* (14.57) gave values that were statistically significantly higher compared to the *third* year (12.17). Leaf area measurement (cm² per plant) showed statistically significant differences among treatments (Table 3). An interaction of magnetic field and variety has been found, in the three-year field experiment (Table 1). The highest leaf area has been found at *Grecale-45* treatment (249.26), with statistically significant differences compared to all other treatments, followed by *Grecale-30* (240.13). This treatment gave values with statistically significant differences compared to *Grecale-15* (230.94), which also gave values with

statistically significant differences compared to *Grecale-0* (219.30). Moreover, *Simeto-15* (181.58), *Simeto-30* (176.61) and *Simeto-45* (174.41) gave values that were statistically significantly higher compared to *Simeto-0* (159.82). Regarding year, the *third* (206.88) and the *second* (205.44) gave values that were statistically significantly higher compared to the *first* year (199.71).

Physiology measurements

The chlorophyll content ($\mu\text{g}/\text{cm}^2$) of durum wheat plants was higher at the magnetic field treatments (Table 4). An interaction of magnetic field and variety has been found, in the three-year field experiment (Table 1). The highest values of chlorophyll content were measured at *Simeto-15* treatment (49.27), *Simeto-30* (48.61), *Simeto-45* (48.51) and *Grecale-30* (47.90) with statistically significant differences compared to all other treatments except *Grecale-45* (47.04). Moreover, *Grecale-15* (45.50) gave values that were no statistically significantly higher compared to *Grecale-45* and *Grecale-0* (43.94). *Grecale-0* gave values that were statistically significantly higher compared to *Simeto-0* (38.95). Regarding year, the *first* (54.06) and the *second* (52.74) gave values that were statistically significant higher compared to the *third* year (54.06).

Photosynthetic rate ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) of durum wheat plants was higher at the magnetic field treatments (Table 5). An interaction of magnetic field and variety has been found, in the three-year field experiment (Table 1). The highest photosynthetic rate has been found at *Simeto-30* (19.52) and *Simeto-45* (19.31) treatments, with statistically significant differences compared to all other treatments, except *Simeto-15* (17.78) and *Grecale-30* treatment (16.98). *Simeto-15* and *Grecale-30* gave values with no statistically significant differences compared to *Grecale-15* (15.89), *Grecale-45* (16.70) and *Simeto-0* (15.52). However, *Simeto-15* and *Grecale-30* gave values with statistically significant differences compared to *Grecale-0*

(11.66). Regarding year, the *third* (19.14) gave values that were statistically significantly higher compared to the *second* year (17.07), which also gave values that were statistically significantly higher compared to the *first* year (13.79). Transpiration rate ($\text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$) of durum wheat plants showed some statistically significant differences regarding magnetic field treatment, variety and year. Treatments of 30 (4.06) and 45 *minutes* (3.90) of magnetic field exposure gave values that were statistically significantly higher compared to 15 *minutes* (3.38). All magnetic field treatments gave values that were statistically significant higher compared to *control* (2.24). Regarding year, the *third* (3.88) and the *second* (3.41) gave values that were statistically significantly higher compared to the *first* year (2.89). Finally, the variety *Simeto* (3.55) gave values that were statistically significant higher compared to *Grecale* (3.24). Stomatal conductance ($\text{mol m}^{-2} \text{ s}^{-1}$) of durum wheat plants was higher at the magnetic field treatments (Table 5). An interaction of magnetic field and variety has been found, in the three-year field experiment (Table 1). The highest stomatal conductance has been found at *Simeto-30* (0.24) and *Simeto-45* (0.24) treatments, with statistically significant differences compared to *Simeto-15* (0.20) treatment. *Simeto-15* gave values that were statistically significantly higher compared to *Grecale-30* (0.18), *Grecale-45* (0.17), *Grecale-15* (0.17) and *Simeto-0* (0.17). *Grecale-0* (0.13) gave the lowest values of all treatments with statistically significant differences.

Yield

The pre-sowing application of magnetic field increased yield of durum wheat (Table 6). Yield (kg per ha) of durum wheat plants showed some statistically significant differences regarding magnetic field treatment, variety and year. The treatments of 45 (3229.2), 30 (3113.4) and 15 *minutes* (2941.8) of magnetic field exposure gave values that were statistically significantly higher compared to 0 *minutes* (2428.5). That means that all magnetic field treatments gave values that were statistically significantly higher compared to *control*. Regarding year, the

second (3332.8) gave values that were statistically significantly higher compared to the *third* year (3045.5). The *third* year gave values that were statistically significantly higher compared to the *first* year (2406.3). Finally, the variety *Grecale* (3126.7) gave values that were statistically significantly higher compared to *Simeto* (2729.8).

Activity of α -amylase

The activity of α -amylase has been conducted on Simeto variety. The results showed that pre-sowing application of magnetic field on durum wheat seeds leads to increased activity of α -amylase at 30 minutes of exposure, the third day after sowing. Moreover, at the fourth day after sowing, in all magnetic field measurements (15, 30 and 45 minutes) the observed enzyme activity was statistically significantly higher compared to the untreated seeds (*control*). At the fifth day, all treatments gave similar results, with no statistically significant differences.

DISCUSSION

The results obtained in this three-year experiment showed a positive impact of pulsed electromagnetic field, in durum wheat cultivation. Magnetic field has been found to enhance germination, tillering, dry weight, leaf area, chlorophyll content, photosynthetic rate, transpiration rate, stomatal conductance and yield of two durum wheat cultivars. In addition, α -amylase activity measurements showed that magnetic field affects enzymes and this could possibly explain the improved germination.

In 1999 Pietruszewski found that the presowing application of magnetic field increased the germination of two wheat varieties, while similar are the findings of Aksenov in 1997. Increase of the percentage of germination rate as a result of exposure to a magnetic field states also Cakmak (2010) for wheat grain. The positive influence of the magnetic field in the

germination of corn seeds has also been expressed and through the reduction of time of germination. Florez in 2007 indicates that this reduction eventually led to an increase of 16 to 25% germination rate compared to the control. It should be noted that the emergence of the capacity improvement is recorded in controlled conditions (Petri dishes) and field conditions (Bilalis et al., 2012b).

Recently, Javed et al., (2011) found that pretreated corn seeds with different electromagnetic treatments particularly 100 and 150 mT for 10 min significantly alleviated the drought-induced adverse effects on growth by improving photosynthesis, transpiration rate and stomatal conductance. Pulsed Electromagnetic Field (PEMF) improved main physiology measurements such transpiration rate, photosynthetic rate and stomatal conductance at the early stages of cotton plants (Bilalis et al., 2013).

Low-intensity magnetic field has been used in strawberry plants and had a positive effect on the yields, in weight and in the number of fruit (Esitken and Turan, 2004). The pre treatment of lettuce seeds with magnetic field also led to an increase in final yield (De Souza et al., 2008). With presowing application of the magnetic field in cotton, it has been found that certain varieties can double their yields compared to control (Leelapriya et al., 2003). A recent study used two types of hybrid corn using three different durations of exposure to the magnetic field. The results showed that the ideal exposure to the magnetic field varies even among the groups of the same species (Bilalis et al., 2012c). A similar finding has been noted by other researchers who argue that it is necessary to find the exact parameters of radiation that cause positive biostimulation in seeds, which depends on the genotype (Hernandez-Aguilar et al., 2009). Pietruszewski (1996) has reported increase in crop yields after the influence of the magnetic field.

The exposure of seeds at magnetic field was found to increase the activity of α -amylase in the third and fourth day after sowing, thus giving an advantage during germination. On the fourth day after seeding, the magnetic field treatments exhibited increased activity of 50 to 100% compared with the control treatment. Amylases are enzymes that catalyze the hydrolysis of starch in the early stages of seed germination. The hydrolytic enzymes secreted by diktyosomata inside the endosperm and catalyze the hydrolysis of starch to glucose reserves. Glucose is the source of energy and carbon skeletons for growth of the young seedling, which in the early stages of life remains heterotrophic. Recent studies have reported higher activity of α -amylase in sunflower seeds having treated with magnetic field (Vashisth and Nagarajan, 2010). In contrast, in soybean seeds the pretreatment with PMF resulted to α -amylase activity reduction eight days after treatment. (Radhakrishnan and Kumari, 2012).

The multiple regression analysis indicates that there was a statistically significant relationship between Dry Weight (DW150), Leaf Area (LA150) and Yield (YIELD):

$$Yield = 32.024 + 7.048 \times DW150 + 0.8 \times LA150$$

	St. Error	p-level
DW150	3.314	0.037
LA150	0.175	0.000

This equation explains that Leaf Area ($p < 0.001$) and Dry Weight ($p < 0.05$) affected statistically significant the Yield of durum wheat (Figure 2).

In a similar field experiment conducted in tomato crop, magnetic field treatment, in certain times of exposure, improved shoot diameter, number of leaves per plant, fresh and dry weight, number of flowers, and yield per plant.

CONCLUSIONS

Pre-sowing pulsed electromagnetic field treatment enhanced germination, tillering, dry weight, leaf area, chlorophyll content, photosynthetic rate, transpiration rate and stomatal conductance of two durum wheat cultivars. Duration of 30 and 45 minutes of pre-sowing magnetic field treatment gave the best results. The exposure of seeds at magnetic field was found to increase the activity of α -amylase in the third and fourth day after sowing, thus giving an advantage during germination. Enhancement of germination, tillering, plant growth and physiology of durum wheat plants has led to increased yield in this three year experiment. Such studies provide a holistic approach of an agricultural cultivation that can lead to the comprehension of the exact mechanism of magnetic field effect on plant tissues and lead to the appropriate application of magnetic fields (Efthimiadou et al., 2014).

Table 1. Analysis of variance (F values)

	Germination (plants per row) 20 DAS	Tillering (plants per row) 50 DAS	Dry Weight (g per plant) 150 DAS	Leaf Area (cm² per plant) 150 DAS
Year	6.39 ^{ns}	57.26 ^{***}	493.97 ^{***}	9.13 [*]
Variety	348.36 ^{***}	586.31 ^{***}	19.37 [*]	1017.8

					6***
Magnetic	178.08***	71.51***	249.00***	88.10**	*
Variety*Magnetic	18.46***	9.05***	3.63*	13.33**	*
Year*Variety*Magnetic	1.27 ^{ns}	2.75 ^{ns}	4.39 ^{ns}	2.76 ^{ns}	
	Chlorophyll content (µg per cm²)	Photosynthetic rate (µmol CO₂m⁻² s⁻¹)	Transpiration rate (mmol H₂O m⁻² s⁻¹)	Stomatal conductance (mol m⁻² s⁻¹)	Yield (kg per hectare)
	130 DAS	115 DAS	115 DAS	115 DAS	
Year	1399.29***	262.28***	41.17**	5.71 ^{ns}	172.35***
Variety	0.51 ^{ns}	233.72***	17.53*	70.08**	98.26***
Magnetic	77.05***	120.14***	76.94***	129.12***	38.24***
Variety*Magnetic	30.19***	3.63*	0.95 ^{ns}	14.67***	2.63 ^{ns}
Year*Variety*Magnetic	1.60 ^{ns}	1.65 ^{ns}	1.18 ^{ns}	0.81 ^{ns}	1.48 ^{ns}

*, **, ***=significance at 0.05, 0.01 and 0.001. ns=not significant

Table 2. Interaction of magnetic field and variety on germination (30 DAS) and tillering (50 DAS) of durum wheat for three years.

Germination (plants per row) 20 DAS				Tillering (plants per row) 50 DAS			
<i>Magnetic*Variety</i>		<i>Year</i>		<i>Magnetic*Variety</i>		<i>Year</i>	
Simeto 0	60.05 e	First	74.30 a	Simeto 0	78.17 e	First	106.37 a
Simeto 15	69.10 d	Second	74.84 a	Simeto 15	92.04 d	Second	105.81 a

Simeto 30	73.17 c	Third	73.10 a	Simeto 30	95.73 d	Third	100.20 b
Simeto 45	78.34 b			Simeto 45	96.62 d		
Grecale 0	71.68 c			Grecale 0	104.62 c		
Grecale 15	78.80 ab			Grecale 15	112.81 b		
Grecale 30	80.55 ab			Grecale 30	123.86 a		
Grecale 45	80.96 a			Grecale 45	129.16 a		

Means followed by the same letter for treatments are not significantly different according to the least significant difference (LSD) test

Table 3. Interaction of magnetic field and variety on dry weight and leaf area (150 DAS) of durum wheat for three years.

Dry Weight (g per plant) 150 DAS				Leaf Area (cm ² per plant) 150 DAS			
<i>Magnetic*Variety</i>		<i>Year</i>		<i>Magnetic*Variety</i>		<i>Year</i>	
Simeto 0	12.03 c	First	14.57 a	Simeto 0	159.82 f	First	199.71 b
Simeto 15	13.81 b	Second	14.82 a	Simeto 15	181.58 e	Second	205.44 a
Simeto 30	14.05 ab	Third	12.17 b	Simeto 30	176.61 e	Third	206.88 a
Simeto 45	15.23 a			Simeto 45	174.41 e		
Grecale 0	11.94 c			Grecale 0	219.30 d		
Grecale 15	14.02 ab			Grecale 15	230.94 c		
Grecale 30	14.64 ab			Grecale 30	240.13 b		
Grecale 45	15.11 ab			Grecale 45	249.26 a		

Means followed by the same letter for treatments are not significantly different according to the LSD test.

Table 4. Interaction of magnetic field and variety on chlorophyll (130 DAS) of durum wheat for three years.

Chlorophyll content ($\mu\text{g per cm}^2$) 130 DAS

<i>Magnetic*Variety</i>		<i>Year</i>	
Simeto 0	38.95 d	First	54.06 a
Simeto 15	49.27 a	Second	52.74 a
Simeto 30	48.61 a	Third	31.85 b
Simeto 45	48.51 a		
Grecale 0	43.94 c		
Grecale 15	45.50 bc		
Grecale 30	47.90 a		
Grecale 45	47.04 ab		

Means followed by the same letter for treatments are not significantly different according to the LSD test.

Table 5. Interaction of magnetic field and variety on photosynthetic rate (115 DAS), effect of magnetic field, variety and year on transpiration rate (115 DAS) and interaction of magnetic field and variety on stomatal conductance (115 DAS) of durum wheat for three years. Means followed by the same letter for treatments are not significantly different according to the LSD test.

Photosynthetic rate ($\mu\text{mol CO}_2\text{m}^{-2}\text{s}^{-1}$) 115 DAS		Transpiration rate ($\text{mmol H}_2\text{O m}^{-2}\text{s}^{-1}$) 115 DAS			Stomatal conductance ($\text{mol m}^{-2}\text{s}^{-1}$) 115 DAS	
<i>Magnetic*Variet</i>	<i>Year</i>	<i>Magne</i>	<i>Year</i>	<i>Vari</i>	<i>Magnetic</i>	<i>Year</i>

<i>y</i>		<i>tic</i>				<i>ariety</i>							
Simeto	15.52	First	13.79	2.2	First	2.	Sim	3.	Sim	0.17	First	0.	
0	b			0									
		Seco	17.07		Sec	3.	Grec	3.			Seco	0.	
		nd	b		ond	41	ale	24	Sim		nd	1	
Simeto	17.78			1 3.3		a		b	eto	0.20		8	
15	ab			5 8 b					15	b		a	
		Thir	19.14		Thir	3.					Thir	0.	
		d	a		d	88			Sim		d	1	
Simeto	19.52			3 4.0		a			eto	0.24		9	
30	a			0 6 a					30	a		a	
									Sim				
Simeto	19.31			4 3.9					eto	0.24			
45	a			5 0 a					45	a			
Grecale	11.66								Grec	0.13			
0	c								ale 0	d			
									Grec				
Grecale	15.89								ale	0.17			
15	b								15	c			
									Grec				
Grecale	16.98								ale	0.18			
30	ab								30	c			
									Grec				
Grecale	16.70								ale	0.17			
45	b								45	c			

Table 6. Effect of magnetic field, variety and year on yield of durum wheat. Means followed by the same letter for treatments are not significantly different according to the LSD test.

Yield					
<i>Magnetic</i>		<i>Year</i>		<i>Variety</i>	
0	2428.5 b	First	2406.3 c	Simeto	2729.8 b
15	2941.8 a	Second	3332.8 a	Grecale	3126.7 a
30	3113.4 a	Third	3045.5 b		
45	3229.2 a				

Figure 1. Effect of magnetic field on α -amylase activity. Means followed by the same letter are not significantly different according to the LSD test.





Figure 2. Multiple regression analysis of Yield, Dry Weight and Leaf Area.



References

- Aguilar, C.H., Pacheco A.D., Carballo A.C., Orea A.C., Ivanov R., Bonilla J.L.L., Montanez J.P.V., 2009. Alternative magnetic field irradiation effects on three genotype Maize seed field performance. *Acta Agrophys.* 14(1), 7-17.

- Athanasiou, A., Karkambounas, S., Batistatou, A., Lykoudis, E., Katsaraki, A., Kartsiouni, T., Papalois, A., Evangelou, A., 2007. The effect of pulsed electromagnetic fields on secondary skin wound healing: An experimental study. *Bioelectromagnetics*. 28, 362-368.
- Bhardwaj J., Anand A., Nagarajan S., 2012. Biochemical and biophysical changes associated with magnetopriming in germinating cucumber seeds. *Plant Physiol Biochem*. 57, 67-73.
- Bhatnagar, D., Deb, A.R., 1978. Some aspects of pregermination exposure of wheat seeds to magnetic field II. Effect on some physiological processes. *Seed Res*. 6, 14-22.
- Bilalis, D., Katsenios, N., Efthimiadou, A., Karkanis A., 2012a. Pulsed electromagnetic fields effect in oregano rooting and vegetative propagation: A potential new organic method. *Acta Agr. Scand. B-S P*. 62, 94-99.
- Bilalis, D., Katsenios, N., Efthimiadou, A., Karkanis A., 2012b. Investigation of pulsed electromagnetic field as a novel organic pre-sowing method on germination and initial growth stages of cotton. *Electromagn. Biol. Med*. 31(2), 143-150.
- Bilalis, D., Katsenios, N., Efthimiadou, A., Karkanis A., 2012c. Pulsed electromagnetic field: an organic compatible method to promote plant growth and yield in two corn types. *Electromagn. Biol. Med*. 31(4), 333-343.
- Bilalis, D., Katsenios, N., Efthimiadou, A., Karkanis A. Khah M.E., Mitsis T., 2013. Magnetic field pre-sowing treatment as an organic friendly technique to promote plant growth and chemical elements accumulation in early stages of cotton. *Aust. J. Crop Sci*. 7(1), 46-50.

- Cakmak, T., Dumlupinar, R., Erdal, S., 2010. Acceleration of germination and early growth of wheat and bean seedlings grown under various magnetic field and osmotic conditions. *Bioelectromagnetics*. 31, 120-129.
- Çelik, Ö., Atak Ç., Rzakoulieva A., 2008. Stimulation of rapid regeneration by a magnetic field in Paulownia node cultures. *J Cent Eur Agr*. 9(2), 297-304.
- Dardeniz A., S. Tayyar, S. Yalcin, 2006. Influence of Low-Frequency Electromagnetic Field on the Vegetative Growth of Grape cv. Uslu. *J Cent Eur Agr*. 7(3), 389-395.
- De Souza , A., García D., Sueiro L., Gilart F., Porras E., Licea L., 2006. Pre-sowing magnetic treatments of tomato seeds increase the growth and yield of plants. *Bioelectromagnetics*. 27(4), 247-257.
- De Souza, A., Sueiro, L., Gonzalez, L. M., Licea L., Porras E. P., Gilart., 2008. Improvement of the growth and yield of lettuce plants by non-uniform magnetic fields. *Electromagn. Biol. Med*. 27, 173-184.
- De Souza, A., García D, Sueiro L, Gilart F., 2014. Improvement of the seed germination, growth and yield of onion plants by extremely low frequency non-uniform magnetic fields. *Sci Hortic-Amsterdam*. 176, 63–69.
- Dhawi, F.& Al-Khayri, J.M., 2009. The effect of magnetic resonance imaging on date palm (*Phoenix dactylifera* L.) elemental composition. *Communications in Biometry and Crop Science*. 4(1), 14–20.
- Efthimiadou A., Katsenios N., Karkanis A., Papastilianou P. ,Triantafyllidis V., Travlos I. Bilalis J.D., 2014. Effects of Presowing Pulsed Electromagnetic Treatment of Tomato Seed on Growth, Yield, and Lycopene Content. *The Scientific World Journal*. 369745, 6 pages.

- Esitken, A., & Turan, M., 2004. Alternating magnetic field effects on yield and plant nutrient element composition of strawberry (*Fragaria ananassa* cv. *camarosa*). *Acta Agr. Scand. B-S P.* 54, 135-139.
- Flórez, M., Carbonell, M.V. Martínez, E. 2007. Exposure of maize seeds to stationary magnetic fields: Effects on germination and early growth. *Environ Exp Bot.* 59(1), 68-75.
- Hajnorouzi, A., Vaezzadeh, M., Ghanati F. , 2011. Growth promotion and a decrease of oxidative stress in maize seedlings by a combination of geomagnetic and weak electromagnetic fields. *J Plant Physiol.* 168(10), 1123-1128.
- Javed N, Ashraf M, Akram NA, Al-Qurainy F., 2011. Alleviation of Adverse Effects of Drought Stress on Growth and Some Potential Physiological Attributes in Maize (*Zea mays* L.) by Seed Electromagnetic Treatment. *Photochem Photobiol.* 87, 1354-1362.
- Katsenios, N., 2013. Investigation of pulsed electromagnetic field influence on plant growth. Ph.D. Thesis. Agricultural University of Athens, Greece.
- Leelapriya T., Dhilip K.S. and Sanker Narayan P.V., 2003. Effect of weak sinusoidal magnetic field on germination and yield of cotton (*Gossypium* spp.). *Electromag. Biol. Med.* 22, 117-125.
- Lichtenthaler, HK and Wellburn AR., 1983. Determinations of total carotenoids and chlorophylls *a* and *b* of leaf extracts in different solvents. *Biochem Soc Trans.* 11, 591-592.
- Martínez, E., Flórez, M., Maqueda, R., Carbonell, M.V., Amaya, J.M., 2009. Pea (*Pisum sativum*, L.) and lentil (*Lens culinaris*, Med.) growth stimulation due to exposure to 125 and 250 mT stationary fields. *Pol J Environ Stud.* 18(4), 657-663.

- Milgram J, Shahar R, Levin-Harrus T, Kass P., 2004. The effect of short high intensity magnetic field pulses on the healing of skin wounds in rat. *Bioelectromagnetics*. 25, 271-277.
- Muraji M., Asai T., Tatebe W., 1998. Primary root growth rate of Zea mays seedlings grown in an alternating magnetic field of different frequencies. *Bioelectroch Bioener* 44, 271-273.
- Muszynski S., Gagos M. Pietruszewski S., 2009. Short-term pre-germination exposure to ELF magnetic fields does not Influence seedling growth in durum wheat (*Triticum durum*). *Pol J Environ Stud*. 18(6), 1065-1072.
- Odhiambo, J.O., Ndiritu F.G. & Wagara I.N., 2009. Effects of Static Electromagnetic fields at 24 hours incubation on the germination of Rose Coco Beans (*Phaseolus vulgaris*) *Rom J Biophys*. 19(2), 135-147.
- Pietruszewski S., 1996). Effects of magnetic biostimulation of wheat seeds on germination, yield and proteins. *Int. Agrophysics*. 10(1), 51-55.
- Podlesny J., Pietruszewski S., Podlesna A., 2004. Efficiency of the magnetic treatment of broad bean seeds cultivated under experimental plot conditions. *Int. Agrophysics*. 18, 65-71.
- Podlesny J., Pietruszewski S., Podlesna A., 2005. Influence of magnetic stimulation of seeds on the formation of morphological features and yielding of the pea. *Int. Agrophysics*. 19, 61-68.
- Poinapen D, Brown DC, Beeharay GK., 2013. Seed orientation and magnetic field strength have more influence on tomato seed performance than relative humidity and duration

of exposure to non-uniform static magnetic fields. J Plant Physiol. 170(14), 1251-1258.

Stavroulakis, P., 2003. Biological Effects of Electromagnetic Fields. Mechanisms, Modeling, Biological Effects, Therapeutic Effects, International Standards, Exposure Criteria, XVI, 793 p.280.

Torres, C., Diaz, J. E. & Cabal, P. A., 2008. Magnetic fields effect over seeds germination of rice (*Oryza sativa* L.) and tomato (*Solanum lycopersicum* L.). Agronomia Colombeana, vol.26, No.2, pp.177-185.

Vashisth A., Nagarajan S., 2008. Exposure of seeds to static magnetic field enhances germination and early growth characteristics in chickpea (*Cicer arietinum* L.). Bioelectromagnetics. 29(7), 571-578.

Vashisth, A., Nagarajan, S., 2010. Effect on germination and early growth characteristics in sunflower (*Helianthus annuus*) seeds exposed to static magnetic field. J Plant Physiol. 167, 149-156.

Vashisth A., Meena N., Joshi D, K., Singh R., 2013. Study of biophysical and growth parameters in relation to yield in sunflower (*Helianthus annuus*). Indian J Agr Sci. 83 (2).